

2217-913

5/7/2014

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

James L. Kuntsman, Ph.D.
PBI/Gordon Corporation
1217 W. 12th Street
P.O. Box 014090
Kansas City, Missouri 64101

MAY -7 2014

Subject: Notification; Per PR-Notice 98-10
EH-1451 Weed & Feed
EPA Reg. No. 2217-913
Date Submitted: May 6, 2014

Dear Dr. Kuntsman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated May 6, 2014 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact Kable Bo Davis at (703) 306-0415 or davis.kable@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be "K. Montague", written over a circular stamp or seal.

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

for



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 2217-913	2. EPA Product Manager Kathryn V. Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) EH-1451 Weed & Feed	PM# Product Manager—Team 23	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAY - 7 2014
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments, Section(s) II(N) and IV(A) and (E). Please refer to cover letter for details.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

e-mail to jkunstman@pbigordon.com FAX: 816-421-2731

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per Container	<input type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 16-50 pound bags		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name James L. Kunstman, Ph.D.		Title Director of Regulatory Services		Telephone No. (Include Area Code) 816-460-6292	
<p>Certification</p> <p><i>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</i></p>					6. Date Application Received (Stamped)
2. Signature 		3. Title Director of Regulatory Services			
4. Typed Name James L. Kunstman, Ph.D.		5. Date May 6, 2014			

#112

EH-1451 WEED AND FEED

NOTIFICATION

MAY - 7 2014

EPA Reg. No. 2217-913

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	1.108%
Mecoprop-p	0.167%
Dicamba	0.071%
Dithiopyr	0.160%

OTHER INGREDIENTS:	98.494%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4)
 - 3.34 lb (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167% (CAS 16484-77-8)
 - 1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9)
 - 3.20 lb dithiopyr per ton or 0.160% (CAS 97886-45-8)
- Isomer Specific By AOAC Methods.

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen (N)	x.x%
x.x% Ammoniacal Nitrogen	
x.x% Water Insoluble Nitrogen	
x.x% Urea Nitrogen	
x.x% Other Water Soluble Nitrogen	
Available Phosphate (P2O5)	x.x%
Soluble Potash (K2O)	x.x%
Iron (Fe)	x.x%
Nutrient Sources: [Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer.]	

KEEP OUT OF REACH OF CHILDREN

CAUTION

[For residential use by homeowners:]
New York State: Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year [0.25 lb. of dithiopyr per acre per year].

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.

PRODUCT DESCRIPTION

EH-1451 Weed & Feed is formulated (designed) to provide control of most annual grasses and certain broadleaf weeds commonly found in home lawns.

EH-1451 Weed & Feed contains three components: a preemergent herbicide, a lawn fertilizer, and a post emergent broadleaf weed killer.

- The preemergent herbicide can be applied to established turfgrass to control crabgrass goosegrass, foxtails, and other undesirable annual grasses.
- The fertilizer contains slow release nutrients that may extend the availability for turfgrass uptake.

- The post emergent broadleaf herbicide will control over 250 lawn weeds including dandelion, chickweed, thistle and clover. A partial list of broadleaf weeds is presented on this label.

WHERE TO USE

EH-1451 Weed & Feed may be used on the following established turf species:

Established Cool Season Turfgrass:

- Kentucky bluegrass
- perennial ryegrass
- fine fescues
- tall fescue

Established Warm Season Turfgrass:

- bahiagrass
- bermudagrass
- buffalograss
- zoysiagrass

Do not use EH-1451 Weed & Feed on dichondra, carpetgrass or colonial bentgrass. Do not use EH-1451 Weed & Feed on St. Augustinegrass in Florida. Do not use this product on turfgrass where desirable clovers are present.

Guidelines to follow:

The use of EH-1451 Weed & Feed on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

Other Use Precautions:

- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.
- Do not apply when windy.
- Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.
- Do not exceed specified dosage for any area.
- Do not use clippings from treated areas for composting or for mulching vegetables or fruit trees.

WEEDS CONTROLLED			
When applied as directed, this product will control and/or suppress* the following weeds:			
Annual Grass Weeds:			
Annual bluegrass* (<i>Poa annua</i>)	Annual ryegrass Barnyardgrass*	Crabgrass Foxtail (green and yellow)	Goosegrass* (crowfootgrass)
A partial listing of established broadleaf weeds: [see appendix for complete list or include here]			
Bedstraw	Curly dock	Mustard	Red sorrel
Beggarticks	Daisy, oxeye	Narrowleaf plantain	Redroot pigweed
Black medic	Dandelion	Oxalis	Sheep sorrel
Broadleaf plantain	Dichondra	Parsley-piert	Shepherdspurse
Buckhorn plantain	False dandelion	Pearlwort	Slender plantain
Burclover	Ground ivy	Pennycress	Smooth dock
Burdock	Healall	Pepperweed	Smooth pigweed
Carpetweed	Henbit	Pigweed	Spurge
Chickweed, common	Knotweed	Plantain	Thistle
Chickweed, mouseear	Lambsquarters	Poison ivy	Woodsorrel
Chicory	Lespedeza	Poison oak	Yarrow

Clover, hop Clover, strawberry Clover, white	Mallow Matchweed Morningglory	Pokeweed Purslane, common Ragweed	and other broadleaf weeds
* for acceptable control of barnyardgrass and goosegrass, two applications made at a 6 to 10 week interval are needed			

WHEN TO APPLY

A. Established lawns:


This product may be applied as a single application or as split applications in the spring, early summer, or fall.

For the best results in the spring, apply this product one to two (1 to 2) weeks before the expected germination of crabgrass. When the period of germination for crabgrass and other weeds listed on this label extends into early summer, then a second application can be made at 6 to 10 weeks after the initial application.

For crabgrass and other listed grasses: This product is a preemergent herbicide to control crabgrass (smooth, southern and large) and other susceptible annual grasses. This product may also provide early post emergent control of crabgrass. However, post emergent crabgrass control will be obtained only when this product is applied before the first tiller stage of crabgrass.

[start optional growth stage description]

The graphics below indicate the growth stages of crabgrass, and the optimum timing for application of this product to assure control:



BEST Grassy weed control	GOOD Grassy weed control	POOR Grassy weed control
1 to 2 weeks before germination	Stages 9, 10, 11, or 12 above	Stage 13 or greater

[end optional growth stage description]

For broadleaf weed control:

Best results are obtained during the spring and fall when broadleaf weeds are present, young and actively growing under adequate soil moisture and favorable temperatures (50° to 80°F).

HOW TO APPLY

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Avoid spreading granules onto flowers, vegetables, and ornamental shrubbery or desirable plants.

HOW MUCH TO USE

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass:

Apply 2.5 to 3.6 pounds of product per 1,000 square feet of established turfgrass.

For hybrid bermudagrass, bahiagrass and buffalograss:

Apply 2.0 to 3.0 pounds of product per 1,000 square feet of established turfgrass.

[For residential use by homeowners:]

New York State: Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year [0.25 lb. active ingredient of dithiopyr per acre per year].

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application [1.15 lb 2,4-D ae, 0.26 lb MCPP-p acid equivalent, 0.11 lb dicamba acid equivalent, and 0.25 lb dithiopyr per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft. [2.30 lb 2,4-D acid equivalent, 0.52 lb MCPP-p acid equivalent, 0.22 lb dicamba acid equivalent, and 0.50 lb. dithiopyr per acre].

SUGGESTED SPREADER SETTINGS

Note to reader: A listing of spreaders and suggested settings for the application rate of 2.0 to 3.6 lb per 1,000 sq.ft. will be presented on the container labels.

The above spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

CALIBRATION INSTRUCTIONS [examples]:

- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds or other pesticides.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS

• [Controls][Kills][253][more than ###] weeds! (List of weeds may include the following):

Amaranth, purple	Cupid shaving Brush	Mustard, ball	Speedwell, common
Amaranth, slender	Cudweed	Mustard, black	Speedwell, corn
Aster, heath	Daisy, English	Mustard, blue	Speedwell, ivyleaf
Aster, white heath	Daisy, oxeye	Mustard, hedge	Speedwell, persian
Aster, purple	Dandelion, common	Mustard, indian,	Speedwell, purslane
Aster, white prairie		Mustard, tansy	Speedwell, slender
Artichoke	Day flower	Mustard, tumble	Speedwell, snow
Austrian fieldcress	Deadnettle	Mustard, white	Speedwell, thymeleaf
Bedstraw, Catchweed	Dichondra	Mustard, wormseed	Spiny amaranth
Bedstraw, smooth	Dock, broadleaf	Nettle, stinging (burning)	Spurge, spotted
Bedstraw, Northern	Dock, curly	Nettle, tall	Spurge, prostrate
Bedstraw, pineywoods	Dogbane	Nettle, wood	Spurweed
Beggarticks, hairy	Dogfennel	Parsley-piert	Star of Bethlehem
Betony, Florida	Dollarweed	Parsnip	Strawberry, India mock
Bindweed, Field	(Pennywort)	Pearlwort	Tansy ragwort
Bindweed, Hedge	Dove weed	Pennycress (frenchweed)	Tanweed
Bitter Wintercress	Elderberry	Pepperweed, perennial	Thistle, blessed
Bittercress, hairy	Falseflax	Pepperweed, virginia	Thistle, bull
Bitterweed	False sunflower	(Peppergrass)	Thistle, canada
Black-eyed Susan	Fiddleneck	Pigweed, prostrate	Thistle, flodman (prairie)
Black Medic	Filaree, redstem	Pigweed, redroot	Thistle, musk
Blood flower milkweed	Filaree, whitestem	Pigweed, russian	Thistle, platte
Blue lettuce	Fleabane, daisy	Pigweed, smooth	Thistle, plumeless
Brassbuttons	(annual)	Pigweed, tumble	Thistle, russian
Bristly oxtongue	Fleabane, rough	Pineapple weed	Thistle, scotch
Broomweed	Galinsoga, hairy	Plains coreopsis (tickseed)	Thistle, wavyleaf
Buckhorn	Galinsoga, smallflower	Plantain, blackseed	Thistle, yellow (horrible)
Bullnettle	Goldenrod	Plantain, bracted	Trailing crownvetch
Burclover	Ground ivy	Plantain, broadleaf	Velvetleaf
Burdock	Groundsel	Plantain, buckhorn	Vervain, blue
Burweed, lawn (spurweed)	Gumweed	Plantain, hoary	Vervain, hoary
Buttercup, bulbous	Hairy fleabane	Plantain, narrowleaf	Vervain, prostrate
Buttercup, creeping	Hawkweed, meadow	Plantain, slender	Vervain, tall
Buttonweed, common	Hawkweed, mouseear	Plantain, woolly	Vetch, bird
(poorjoe)	Hawkweed, orange	Poison ivy	Vetch, hairy
Buttonweed, virginia	Hawkweed, yellow	Poison oak	Vetch, milk
Carolina geranium (wild)	Healall	Pokeweed	Vetch, narrowleaf
Carpetweed	Heartleaf drymary	Prairie Sunflower	Vetch, wild
Catnip	Hemp	Prickly lettuce (compass plant)	Virginia creeper
Catsear, spotted (false dandelion)	Henbit	Prickly sida	Western clematis
Chamber bitter	Hoary cress	Prostrate knotweed	Western salsify
Chickweed, common	Horsenettle	Puncturevine	Wild aster
Chickweed, mouseear	Horseweed	Purslane, common	Wild buckwheat
Chickweed, sticky	Jimsonweed	Pusley, brazil	Wild carrot
Chicory	Johnny-Jumpup violet	Pusley, florida	Wild four-o'clock
Cinquefoil	Knawel	Pusley, large flower	Wild garlic
Clover, crimson	Knotweed, prostrate	Ragweed, bur	Wild lettuce
Clover, alyce	Kochia	Ragweed, common	Wild marigold
Clover, hop	Lambsquarters	Ragweed, lancleaf	Wild mustard (yellow rocket)
Clover, rabbit foot	Lespedeza	Ragweed, western	Wild onion
Clover, red	Mallow, alkali	Redstem filaree	Wild parsnip
Clover, strawberry	Mallow, bristly	Rough cinquefoil	Wild radish
Clover, yellow sweet	Mallow, common	Scarlet pimpernel	Wild rape
Clover, white sweet	Mallow, dwarf	Shepardspurge	Wild strawberry
Clover, white	Mallow, venice	Smartweed, ladythumb	Wild sweet potato
Cockle	Marcela	Smartweed, pale	Wild violet
Cocklebur	Matchweed	Smartweed, pennsylvania	Woodsorrel, creeping
Cocklebur, oriental	Mexicanweed	Smooth chaff-flower	Woodsorrel, violet
Cocklebur, spiny	Moneywort	Smoothdock	Woodsorrel, yellow (Cxalis)
	Morningglory, bigroot	Sorrel, red (sheep)	Woolly croton

11/12

Common mullein	Morningglory, ivyleaf	Sowthistle, annual	Wormseed
Corn Spurry	Morningglory, mexican	Sowthistle, spiny	Yarrow, common
Creeping Beggarweed	Morningglory, red	Spanish needles	Yarrow, western
Creeping Jenny (creeping charlie)	Morningglory, tall	Speedwell, bi-lobed	Yarrow, yellow
	Morningglory, wolly		
	Mugwort		

Fertilizer Benefit Claims:

- Greens in 72 Hours & Promotes Thick, Green Growth
- Contains Composite Timed-Release (CTR) Nitrogen for Extended Feeding!
- Encourages Quick Greening & Provides Extended Nitrogen Feeding up to 2 Months
- (Product Name) is specially formulated to complement the phosphorus already present in some soils and give your turf the additional primary nutrients needed for a good turf feeding. A soil test can help you determine if your soil has an adequate supply of phosphorus and if this is the best product for your soil type. Proper watering and mowing practices along with regular feedings will help promote a beautiful green lawn.

General Marketing Claims:

- Northern and Southern Grasses Bluegrass, Fescue, Zoysia, Bermuda, Bahia, Perennial Ryegrass
- Treats Northern and Southern Grasses
- Use on Lawns to Kill the Toughest Weeds like Clover and dandelion.
- Starts working on contact
- [cool][all] weather performance
- Proven [all] [cool] weather performance
- For home lawn care
- For home lawns only
- For use on lawns
- Guaranteed [Results] or Your Money Back*
 - If you are not satisfied with this product for any reason, send your written request for a refund to the address at left, together with the original dated store receipt showing the price you paid. Please allow 6 to 8 weeks for processing.
- Kills Weeds While Feeding Your Lawn
- For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass
- Sta-Green Turf Note – Control Those Pesky Weeds
 - Many broadleaf weeds (dandelion, chickweed, henbit) germinate in the fall but are not visible in your lawn until spring. Applying (product name) in the fall can provide effective results. If needed, you can make a follow-up application in the spring.
- Project Guide – For a thicker, greener, more beautiful lawn, follow (Product Program). It's easy to follow and can be started at anytime by simply applying the product that corresponds with the current season!
- Prevents Crabgrass and Fertilizes
- Weed 'n Feed Plus Crabgrass Preventer
- Product Questions? 855-270-4776 [for VPG supplemental distributor product]
- Visit us at: www.fertilome.com
- Covers up to XXXX square feet
- DIMENSION® is a registered trademark of Dow AgroSciences, L.L.C.
- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

3. State mandated fertilizer labeling requirements:

- Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>
- Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing _____

12/12

pounds of nitrogen per 1,000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.

- If the use rate applies less than 1 pound nitrogen per 1,000 sq.ft. per application, the following statement must appear: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq.ft.



--	--	--	--

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00913.20140506.notif-proposed-highlighted

2. **Reason for Issue:** ad claims, fix typos, reorganize appendix

